

# Vascular Model Repository

## Specifications Document



0063\_H\_PULMGLN\_SVD

Legacy Name: 0099\_0001

Model added: 27 Dec 2021

<b>Species</b>	Human
<b>Anatomy</b>	Pulmonary Glenn
<b>Disease</b>	Single Ventricle Defect
<b>Procedure</b>	Glenn

# Clinical Significance and Background

## Pulmonary Glenn

The pulmonary arteries are the blood vessels responsible for transporting deoxygenated blood to the lungs to perform respiration. A normal pulmonary anatomy involves the main pulmonary artery (MPA) trunk leaving the right ventricle of the heart which then branches off into the left and right pulmonary arteries which then continue to fractally branch towards the lungs. However, after the Glenn procedure (an open-heart surgery done on babies with congenital heart defects) the anatomy of the pulmonary system is slightly modified. The main pulmonary artery trunk is not present. Instead, the superior vena cava (SVC) is directly connected to the left and right pulmonary arteries in a T-shaped 3-way intersection configuration. This is done so that deoxygenated blood coming from the upper body (through the SVC) flows directly to the lungs without passing through the heart to allow for the heart to focus solely on pumping oxygenated blood from the lungs to the rest of the body.

The Glenn procedure is very often followed by the Fontan procedure which then further modifies the pulmonary artery anatomy by detaching the inferior vena cava (IVC) and also connecting it directly to the left and right pulmonary arteries in a 4-way intersection configuration.

## Single Ventricle Defect

A single ventricle defect (SVD) is a type of heart defect that a child is born with. It occurs when one of the two pumping chambers in the heart, called ventricles, is not large enough or strong enough to work correctly. In some cases, the chamber might be missing a valve. Single ventricle defects are rare, affecting only about five out of 100,000 newborns. They are also one of the most complex heart problems, usually requiring at least one surgery. There are several types of single ventricle defects which include but are not limited to: tricuspid atresia, hypoplastic left heart syndrome (HLHS), mitral valve atresia (usually associated with HLHS), single left ventricle, double inlet left ventricle (DILV), double outlet right ventricle (DORV), pulmonary atresia with the intact ventricular septum (PA/IVS), and atrioventricular canal defect (AV Canal).

## Glenn

The Glenn procedure is a type of open-heart surgery. Babies who need this surgery typically have it when they are 4 - 6 months old. The Glenn procedure is done for

children who are born with heart problems like hypoplastic left heart syndrome (HLHS), tricuspid atresia, and double outlet right ventricle. Depending on the heart problem, children may need the Norwood procedure before the Glenn surgery.

The Glenn procedure sends blood from the upper body directly to the lungs. This way, the single ventricle only has to pump blood to the body (and not to the lungs), so it does not have to work as hard.

During the Glenn procedure, the surgeon disconnects the superior vena cava (SVC) from the heart and connects it to the pulmonary artery. Now the blood from the upper part of the body flows directly into the pulmonary artery. The pulmonary artery takes the blood to the lungs. If the baby had the Norwood procedure, the surgeon will remove the shunt that was placed then.

## Clinical Data

### General Patient Data

Age (yrs)	4.75
Sex	Male

### Specific Patient Data

CI (L/min/m <sup>2</sup> )	3.2
P SVC MP cath	12
FS LPA MRI	0.2
FS RPA MRI	0.8

## Notes

Paper patient "D". See [paper](#) for more details. See below for information on the image data.

**Image Modality:** MR

**Image Type:** VTI

**Image Source:** TLAB

**Image Manufacturer:** GE MEDICAL SYSTEMS

## Publications

See the following publications which include the featured model for more details:

Troianowski, G., Taylor, C. A., Feinstein, J. A., & Vignon-Clementel, I. E. (2011). Three-dimensional simulations in Glenn patients: clinically based boundary conditions, hemodynamic results and sensitivity to input data. *Journal of biomechanical engineering*, 133(11).  
<http://www.doi.org/10.1115/1.4005377>

# License

Copyright (c) Stanford University, the Regents of the University of California, Open Source Medical Software Corporation, and other parties.

All Rights Reserved.

Permission is hereby granted, free of charge, to any person obtaining a copy of this data to use the data for research and development purposes subject to the following conditions:

The above copyright notice and the README-COPYRIGHT file shall be included in all copies of any portion of this data. Whenever reasonable and possible in publications and presentations when this data is used in whole or part, please include an acknowledgement similar to the following:

"The data used herein was provided in whole or in part with Federal funds from the National Library of Medicine under Grant No. R01LM013120, and the National Heart, Lung, and Blood Institute, National Institutes of Health, Department of Health and Human Services, under Contract No. HHSN268201100035C"

AND/OR

N.M. Wilson, A.K. Ortiz, and A.B. Johnson, "The Vascular Model Repository: A Public Resource of Medical Imaging Data and Blood Flow Simulation Results," J. Med. Devices 7(4), 040923 (Dec 05, 2013) doi:10.1115/1.4025983.

AND/OR

Reference the official website for this data: [www.vascularmodel.com](http://www.vascularmodel.com)

THE DATA IS PROVIDED "AS IS", WITHOUT WARRANTY OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO THE WARRANTIES OF MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE AND NONINFRINGEMENT. IN NO EVENT SHALL THE AUTHORS OR COPYRIGHT HOLDERS BE LIABLE FOR ANY CLAIM, DAMAGES OR OTHER LIABILITY, WHETHER IN AN ACTION OF CONTRACT, TORT OR OTHERWISE, ARISING FROM, OUT OF OR IN CONNECTION WITH THE DATA OR THE USE OR OTHER DEALINGS IN THE DATA.